

POINCARÉ AND MEAN VALUE INEQUALITIES FOR HYPERSURFACES IN RIEMANNIAN MANIFOLDS AND APPLICATIONS*

HILÁRIO ALENCAR[†] AND GREGÓRIO SILVA NETO[‡]

Abstract. In the first part of this paper we prove some new Poincaré inequalities, with explicit constants, for domains of any hypersurface of a Riemannian manifold with sectional curvatures bounded from above. This inequalities involve the first and the second symmetric functions of the eigenvalues of the second fundamental form of such hypersurface. We apply these inequalities to derive some isoperimetric inequalities and to estimate the volume of domains enclosed by compact self-shrinkers in terms of its scalar curvature. In the second part of the paper we prove some mean value inequalities and as consequences we derive some monotonicity results involving the integral of the mean curvature.

Key words. Poincaré inequality, isoperimetric inequality, monotonicity, hypersurfaces, mean curvature, scalar curvature.

AMS subject classifications. 53C21, 53C42.

1. Introduction and main results. The classical Poincaré inequality states that if $\Omega \subset \mathbb{R}^n$ is a bounded, connected, open subset of \mathbb{R}^n and $1 \leq p < \infty$, then there is a constant $C(p, \Omega)$ depending on p and Ω such that for all non-negative $f \in W_0^{1,p}(\Omega)$ the following inequality holds

$$\left(\int_{\Omega} f^p dx \right)^{\frac{1}{p}} \leq C(p, \Omega) \left(\int_{\Omega} |\nabla f|^p dx \right)^{\frac{1}{p}}, \tag{1.1}$$

where dx denotes the Lebesgue measure of \mathbb{R}^m . If $\Omega = B_r(x_0)$ is the open ball of \mathbb{R}^m with center $x_0 \in \mathbb{R}^m$ and radius r , then there is a constant $C(p)$ depending only on p such that

$$\int_{B_r(x_0)} f^p dx \leq C(p)r \int_{B_r(x_0)} |\nabla f|^p dx,$$

see [16, pp. 289-290]. The Poincaré inequality (1.1) also holds for functions $f \in W^{1,p}(\Omega)$ provided $\int_{\Omega} f dx = 0$, where $\Omega \subset \mathbb{R}^n$ is a bounded, connected open subset of \mathbb{R}^n with Lipschitz boundary $\partial\Omega$ and for all $1 \leq p \leq \infty$, i.e., there is a constant $C(p, \Omega)$ depending on p and Ω such that

$$\left(\int_{\Omega} |f|^p dx \right)^{\frac{1}{p}} \leq C(p, \Omega) \left(\int_{\Omega} |\nabla f|^p dx \right)^{\frac{1}{p}}. \tag{1.2}$$

This is the Poincaré-Wirtinger inequality. An interesting question about these inequalities is to know the dependence of the Poincaré constant $C(p, \Omega)$ on the geometry of the domain Ω or to find the best constant $C(p, \Omega)$ for a given domain and a given p , $1 \leq p \leq \infty$. For convex domains $\Omega \subset \mathbb{R}^n$, Payne-Weinberger showed in [27] that,

[†]Instituto de Matemática, Universidade Federal de Alagoas, Maceió, AL, 57072-900, Brazil (hilario@mat.ufal.br). H. Alencar was partially supported by CNPq of Brazil.

[‡]Instituto de Matemática, Universidade Federal de Alagoas, Maceió, AL, 57072-900, Brazil (gregorio@im.ufal.br).

for $p = 2$, the best Poincaré constant in inequality (1.2) is $C(2, \Omega) = \frac{1}{\pi}(\text{diam } \Omega)$, and Acosta-Duran showed in [1, Thm. 3.2] that, for $p = 1$, the best Poincaré constant is $C(1, \Omega) = \frac{1}{2}(\text{diam } \Omega)$, where here $(\text{diam } \Omega)$ denotes the diameter of Ω in \mathbb{R}^n . Moreover, they showed that, for any value of p , $1 \leq p \leq \infty$, the Poincaré constant for convex domains $\Omega \subset \mathbb{R}^n$ depends only on the diameter of Ω . These Poincaré type inequalities can be extended to domains in Riemannian manifolds. For instance, the following result of P. Li and R. Schoen:

THEOREM 1.1 ([22], Thm. 1.1, p. 282). *Let M be a compact Riemannian m -manifold with boundary ∂M , possibly empty, and with Ricci curvature $\text{Ric}_M \geq -(m - 1)k$ for a constant $k \geq 0$. Let $x_0 \in M$ and $r > 0$. If $\partial M = \emptyset$ assume that the diameter of M is greater than or equal to $2r$. If $\partial M \neq \emptyset$, assume that the distance from x_0 to ∂M is at least $5r$. For every Lipschitz function f on $B_r(x_0)$ which vanishes on $\partial B_r(x_0)$ we have the Poincaré inequality*

$$\int_{B_r(x_0)} |f| dM \leq r(1 + \sqrt{kr})^{-1} e^{2m(1+\sqrt{kr})} \int_{B_r(x_0)} |\nabla f| dM.$$

Poincaré inequalities on domains of Riemannian manifolds has been studied extensively by many authors and it plays an important role in Geometry and Analysis, see [7], [20], [21], [23], [24] and [26] for few examples.

Before to state the results of this paper, let us introduce some definitions and notations. Let M^m , $m \geq 2$, be a m -dimensional hypersurface with boundary ∂M , possibly empty, of a Riemannian $(m + 1)$ -manifold \overline{M}^{m+1} . Let k_1, k_2, \dots, k_m be the principal curvatures of M . We define the first and the second symmetric functions associated to the principal curvatures of M by

$$S_1 = \sum_{i=1}^m k_i \text{ e } S_2 = \sum_{i < j}^m k_i k_j. \tag{1.3}$$

These functions have natural geometric meaning. In fact, $S_1 = mH$, where H denotes the mean curvature of M . If \overline{M}^{m+1} has constant sectional curvature κ , then $2S_2 = m(m - 1)(R - \kappa)$, where R is the scalar curvature of M .

If $K_{\overline{M}} = K_{\overline{M}}(x, \Pi_x)$ denotes the sectional curvature of \overline{M} in $x \in M$ relative to the 2- dimensional subspace $\Pi_x \subset T_x \overline{M}$, we define

$$\kappa_0(x) = \inf_{\Pi_x \subset T_x \overline{M}} K_{\overline{M}}(x, \Pi_x).$$

Let $i(\overline{M})$ be the injectivity radius of \overline{M} and let us denote by $(\text{diam } \Omega)$ the diameter of smallest extrinsic ball which contains $\Omega \subset M$.

In this paper we will address Poincaré type inequalities, with explicit constants, on domains of any hypersurface of a Riemannian manifold with sectional curvatures bounded from above.

THEOREM 1.2 (Poincaré type inequality). *Let \overline{M}^{m+1} be a Riemannian $(m + 1)$ -manifold, $m \geq 2$, with sectional curvatures bounded from above by a constant κ . Let M be a hypersurface of \overline{M}^{m+1} , with boundary ∂M , possibly empty, such that $S_1 > 0$ and $S_2 \geq 0$. Let $\Omega \subset M$ be a connected and open domain with compact closure.*

If $\partial M \neq \emptyset$, assume in addition that $\overline{\Omega} \cap \partial M = \emptyset$. If $(\text{diam } \Omega) < 2i(\overline{M})$ then, for every non-negative C^1 -function $f: M \rightarrow \mathbb{R}$, compactly supported in Ω , we have

$$\int_{\Omega} f S_1 dM \leq C(\Omega) \int_{\Omega} \left[|\nabla f| S_1 + \left(\frac{m(\kappa - \kappa_0)}{4} + S_2 \right) f \right] dM, \quad (1.4)$$

where

$$C(\Omega) = C(1, \Omega) = \begin{cases} \frac{1}{m-1} (\text{diam } \Omega) & \text{if } \kappa \leq 0; \\ \frac{2}{\sqrt{\kappa}(m-1)} \tan\left(\frac{\sqrt{\kappa}}{2} (\text{diam } \Omega)\right), & \text{if } \kappa > 0; \end{cases}$$

and we assume $(\text{diam } \Omega) < \frac{\pi}{\sqrt{\kappa}}$ for $\kappa > 0$. Moreover, if $M = \mathbb{S}^m(r)$ is the Euclidean sphere of radius r , $\overline{M} = \mathbb{R}^{m+1}$, and f is a constant function, then the equality holds.

REMARK 1.1. If $\kappa \leq 0$ and \overline{M} is complete and simply connected, then $i(\overline{M}) = \infty$. If $\kappa > 0$ and the sectional curvatures of \overline{M} are pinched between $\frac{1}{4}\kappa$ and κ , then $i(\overline{M}) \geq \frac{\pi}{\sqrt{\kappa}}$, see for example [13], p.276. For these situations the assumption $(\text{diam } \Omega) \leq 2i(\overline{M})$ in Theorem 1.2 is automatically satisfied.

REMARK 1.2. Results in the direction of the Theorem 1.2 for mean curvature are known, see [28, Theorem 3.1 and Theorem 3.3, p. 531-532].

In the particular case when \overline{M}^{m+1} is one of the space forms \mathbb{R}^{m+1} , $\mathbb{H}^{m+1}(\kappa)$, or $\mathbb{S}^{m+1}(\kappa)$, namely, the Euclidean space, the hyperbolic space of curvature $\kappa < 0$, and the Euclidean sphere of curvature $\kappa > 0$, respectively, we have

COROLLARY 1.1. Let M be a hypersurface with boundary ∂M , possibly empty, of \mathbb{R}^{m+1} , $\mathbb{H}^{m+1}(\kappa)$ or $\mathbb{S}^{m+1}(\kappa)$, $m \geq 2$, with mean curvature $H > 0$ and scalar curvature $R \geq \kappa$. Let $\Omega \subset M$ be a connected and open domain with compact closure. If $\partial M \neq \emptyset$, assume in addition that $\overline{\Omega} \cap \partial M = \emptyset$. Then, for every non-negative C^1 -function $f: M \rightarrow \mathbb{R}$, compactly supported in Ω , we have

$$\int_{\Omega} f H dM \leq C(\Omega) \int_{\Omega} \left[|\nabla f| H + \frac{(m-1)}{2} (R - \kappa) f \right] dM. \quad (1.5)$$

Moreover, if $M = \mathbb{S}^m(r)$ is the Euclidean sphere of radius r in \mathbb{R}^{m+1} and f is a constant function, then the equality holds.

Theorem 1.2 has various immediate consequences that we will present below. The next result should be compared with [5, Theorem 28.2.5, p. 210] which states that

If M^m is a compact submanifold with angles of \mathbb{R}^n , $n > m$, with boundary ∂M , possibly empty, then

$$m \text{vol}(M) \leq R_M \left[\text{vol}(\partial M) + m \int_M |H| dM \right],$$

where R_M is the radius of the smallest ball of \mathbb{R}^n which contains M .

As a consequence of Theorem 1.2 we have

COROLLARY 1.2. Let \overline{M}^{m+1} be a Riemannian $(m+1)$ -manifold, $m \geq 2$, with sectional curvatures bounded from above by a constant κ . Let M be a hypersurface of

\overline{M}^{m+1} , with boundary ∂M , possibly empty, such that $S_1 > 0$ and $S_2 \geq 0$. Let $\Omega \subset M$ be a connected and open domain with compact closure. If $\partial M \neq \emptyset$, assume in addition that $\overline{\Omega} \cap \partial M = \emptyset$. If $(\text{diam } \Omega) < 2i(\overline{M})$ and, for $\kappa > 0$, assuming also $(\text{diam } \Omega) < \frac{\pi}{\sqrt{\kappa}}$, then

$$\int_{\Omega} S_1 dM \leq C(\Omega) \left[\int_{\partial\Omega} S_1 dS_{\Omega} + \int_{\Omega} \left(\frac{m(\kappa - \kappa_0)}{4} + S_2 \right) dM \right],$$

where dS_{Ω} denotes the $(m-1)$ -dimensional measure of $\partial\Omega$. Moreover, if M is compact without boundary, then

$$\int_M S_1 dM \leq C(\Omega) \int_M \left(\frac{m(\kappa - \kappa_0)}{4} + S_2 \right) dM.$$

In particular, if \overline{M} has constant sectional curvature κ , then

$$\int_{\Omega} H dM \leq C(\Omega) \left[\int_{\partial\Omega} H dS_{\Omega} + \frac{m-1}{2} \int_{\Omega} (R - \kappa) dM \right].$$

If M is a compact, without boundary, hypersurface of \mathbb{R}^{m+1} , $\mathbb{H}^{m+1}(\kappa)$ or $\mathbb{S}^{m+1}(\kappa)$, the Corollary 1.2 becomes

COROLLARY 1.3. *Let M be a compact, without boundary, hypersurface of \mathbb{R}^{m+1} , $\mathbb{H}^{m+1}(\kappa)$ and $\mathbb{S}^{m+1}(\kappa)$ with mean curvature $H > 0$ and scalar curvature $R \geq \kappa$.*

(i) *If $M \subset \mathbb{R}^{m+1}$ or $M \subset \mathbb{H}^{m+1}(\kappa)$, then*

$$\int_M H dM \leq \frac{1}{2}(\text{diam } M) \int_M (R - \kappa) dM$$

and

$$\text{diam } M \geq \frac{2 \min H}{\max R - \kappa}.$$

In particular, if $m = 2$ and $M^2 \subset \mathbb{R}^3$, then

$$\int_M H dM \leq 2\pi(\text{diam } M).$$

(ii) *If $M \subset \mathbb{S}^{m+1}(\kappa)$ and $(\text{diam } M) \leq \frac{\pi}{\sqrt{\kappa}}$, then*

$$\int_M H dM \leq \frac{1}{\sqrt{\kappa}} \tan \left(\frac{\sqrt{\kappa}}{2}(\text{diam } M) \right) \int_M (R - \kappa) dM.$$

REMARK 1.3. If $M = \mathbb{S}^m(r)$, the round spheres of radius r in the Euclidean space \mathbb{R}^{m+1} , then the inequalities of the item (i) of Corollary 1.3 become equality.

REMARK 1.4. The proof of the item (i) of the Corollary 1.2 follows taking the compactly supported Lipschitz function $f_{\varepsilon} : M \rightarrow \mathbb{R}$, given by

$$f_{\varepsilon}(x) = \begin{cases} 1 & \text{if } x \in \Omega, \text{dist}(x, \partial\Omega) \geq \varepsilon; \\ \frac{\text{dist}(x, \partial\Omega)}{\varepsilon} & \text{if } x \in \Omega, \text{dist}(x, \partial\Omega) \leq \varepsilon; \\ 0 & \text{if } x \notin \Omega; \end{cases}$$

into the Poincaré inequality (1.4) of Theorem 1.2 and by taking $\varepsilon \rightarrow 0$, where $\text{dist}(x, \partial\Omega)$ denotes the intrinsic distance of M of the point $x \in M$ to the boundary ∂M of M .

A hypersurface $M \subset \mathbb{R}^{m+1}$ is called a self-shrinker if its mean curvature H satisfies $H = -\frac{1}{2m}\langle \bar{X}, \eta \rangle$, where \bar{X} is the position vector of \mathbb{R}^{m+1} and η is the normal vector field of M . Self-shrinkers play an important role in the study of mean curvature flow and they have been extensively studied in recent years, see [6], [11], and [12]. Another consequence of our version of Poincaré inequality is the following result.

COROLLARY 1.4. *Let M be a compact, without boundary, self-shrinker of \mathbb{R}^{m+1} , $m \geq 2$, with mean curvature $H > 0$ and scalar curvature $R \geq 0$. Let $K \subset \mathbb{R}^{m+1}$ be the compact domain such that $M = \partial K$. Then*

$$\text{vol}(K) \leq \frac{m}{m+1}(\text{diam } M) \int_M R dM. \tag{1.6}$$

Moreover, if $M = \mathbb{S}^m(\sqrt{2m})$ is the Euclidean sphere of radius $\sqrt{2m}$, then the equality holds.

The proof follows observing that using divergence theorem,

$$\int_M H dM = \int_{\partial K} \frac{1}{2m} \langle \bar{X}, -\eta \rangle dM = \frac{1}{2m} \int_K \text{div}_{\mathbb{R}^{m+1}} \bar{X} dV = \frac{m+1}{2m} \text{vol}(K)$$

and then replacing the identity above into the item (ii) of Corollary 1.2, for $\kappa = \kappa_0 = 0$ and $2S_2 = m(m-1)R$. The Corollary 1.4 can be compared with Corollary 1.2, p. 4718 of [17], which proves an estimate of the volume of a convex domain in the Euclidean space by the integral of the curvatures of its boundary.

Let \bar{M}^{m+1} be a Hadamard manifold, i.e., a complete and simply connected Riemannian manifold with non-positive sectional curvatures. By the Remark 1.1, $i(\bar{M}) = \infty$. If M is a compact hypersurface of \bar{M}^{m+1} , without boundary, then using the well known D. Hoffman and J. Spruck’s Sobolev inequality, see [19, Thm. 2.1, p. 716], we obtain an estimate for the volume of M .

COROLLARY 1.5. *Let \bar{M}^{m+1} , $m \geq 2$, be a Hadamard manifold with sectional curvatures bounded from above by a constant $\kappa \leq 0$. Let M be a compact hypersurface of \bar{M}^{m+1} , without boundary, such that $S_1 > 0$ and $S_2 \geq 0$. Then*

$$\text{vol}(M)^{\frac{m-1}{m}} \leq \frac{2^{m-1}(m+1)^{1+1/m}}{m(m-1)^2\omega_m^{1/m}}(\text{diam } M) \int_M \left(\frac{m(\kappa - \kappa_0)}{4} + S_2 \right) dM, \tag{1.7}$$

where ω_m denotes the volume of the Euclidean unit sphere.

In fact, by using the Hoffman-Spruck-Sobolev inequality

$$\left[\int_M f^{\frac{m}{m-1}} dM \right]^{\frac{m-1}{m}} \leq \frac{2^{m-2}(m+1)^{1+1/m}}{(m-1)\omega_m^{1/m}} \int_M [|\nabla f| + f|H|] dM$$

for $f \equiv 1$, in addition with the Corollary 1.2, we have

$$\begin{aligned} \text{vol}(M)^{\frac{m}{m-1}} &\leq \frac{2^{m-2}(m+1)^{1+1/m}}{m(m-1)\omega_m^{1/m}} \int_M S_1 dM \\ &\leq \frac{2^{m-1}(m+1)^{1+1/m}}{m(m-1)^2\omega_m^{1/m}}(\text{diam } M) \int_M \left(\frac{m(\kappa - \kappa_0)}{4} + S_2 \right) dM. \end{aligned}$$

1.1. Mean value inequalities. The techniques used in the proof of the Poincaré inequalities can be extended to prove some mean value inequalities involving both S_1 and S_2 . Results in this direction are known for hypersurfaces of the Euclidean space. In this case, holds

THEOREM 1.3 ([8], Lemma 1.18). *Let $M \subset \mathbb{R}^{m+1}$ be a properly immersed hypersurface with mean curvature H and f is a non-negative function on M , then for $s < t$,*

$$\frac{1}{t^m} \int_{M \cap B_t} f dM - \frac{1}{s^m} \int_{M \cap B_s} f dM \geq \int_s^t \frac{1}{r^{m+1}} \int_{M \cap B_r} \langle \bar{X}, \nabla f + fH\eta \rangle dM dr, \tag{1.8}$$

where \bar{X} is the position vector of \mathbb{R}^{m+1} and η is the normal vector field of M .

In the same spirit of inequality (1.8), we obtain

THEOREM 1.4 (Mean Value Inequalities). *Let \bar{M}^{m+1} , $m \geq 2$, be a Riemannian $(m + 1)$ -manifold with sectional curvatures bounded from above by a constant κ . Let M be a proper hypersurface of \bar{M} with $S_1 > 0$ and $S_2 \geq 0$. Let x_0 be a point of \bar{M}^{m+1} , $\rho(x) = \rho(x_0, x)$ be the extrinsic distance function starting at x_0 to $x \in M$, and $B_r = B_r(x_0)$ be the extrinsic ball of center x_0 and radius r . If $\partial M \neq \emptyset$, assume that $B_r \cap \partial M = \emptyset$. Then, for any non-negative, locally integrable, C^1 -function $f : M \rightarrow \mathbb{R}$ and for any $0 < s < t < i(\bar{M})$, we have*

(i) for $\kappa \leq 0$,

$$\begin{aligned} & \frac{1}{t^{\frac{m-1}{2}}} \int_{M \cap B_t} f S_1 dM - \frac{1}{s^{\frac{m-1}{2}}} \int_{M \cap B_s} f S_1 dM \\ & \geq \frac{1}{2} \int_s^t \frac{1}{r^{\frac{m+1}{2}}} \int_{M \cap B_r} \rho [\langle \bar{\nabla} \rho, (S_1 I - A)(\nabla f) + 2S_2 f \eta \rangle + f \text{Ric}_{\bar{M}}(\nabla \rho, \eta)] dM dr; \end{aligned}$$

(ii) for $\kappa > 0$,

$$\begin{aligned} & \frac{1}{(\sin \sqrt{\kappa} t)^{\frac{m-1}{2}}} \int_{M \cap B_t} f S_1 dM - \frac{1}{(\sin \sqrt{\kappa} s)^{\frac{m-1}{2}}} \int_{M \cap B_s} f S_1 dM \\ & \geq \frac{1}{2} \int_s^t \frac{1}{s(\sin \sqrt{\kappa} r)^{\frac{m+1}{2}}} \int_{M \cap B_r} (\sin \sqrt{\kappa} \rho) [\langle \bar{\nabla} \rho, (S_1 I - A)(\nabla f) + 2S_2 f \eta \rangle + f \text{Ric}_{\bar{M}}(\nabla \rho, \eta)] dM dr, \end{aligned}$$

provided $t < \frac{\pi}{2\sqrt{\kappa}}$.

Here $\text{Ric}_{\bar{M}}$ denotes the Ricci tensor of \bar{M} , $A : TM \rightarrow TM$ denotes the linear operator associated with the second fundamental form of M , defined in the tangent bundle TM of M , and $I : TM \rightarrow TM$ denotes the identity operator.

We apply the mean value inequalities of the Theorem 1.4 to obtain some monotonicity results involving the integral of the mean curvature. Monotonicity results appear in several branches of Analysis and Riemannian Geometry in the study to determine the variational behaviour of geometric quantities, see for example [4], [8], [9], [10], [14], [15], [18], [25], [29], and [30].

COROLLARY 1.6 (Monotonicity). *Let \bar{M}^{m+1} , $m \geq 2$, be a Riemannian $(m + 1)$ -manifold with sectional curvatures bounded from above by a constant κ . Let M be a proper hypersurface of \bar{M} with $S_1 > 0$ and $S_2 \geq 0$. If there exists $0 < \alpha \leq 1$, $\Lambda \geq 0$*

and $0 < R_0 < i(\overline{M})$ such that

$$\alpha^{-1} \int_{M \cap B_r} \left(\frac{m(\kappa - \kappa_0)}{4} + S_2 \right) dM \leq \Lambda \left(\frac{r}{R_0} \right)^{\alpha-1} \int_{M \cap B_r} S_1 dM, \quad (1.9)$$

for all $r \in (0, R_0)$, then

(i) for $\kappa \leq 0$, the function $h : (0, R_0) \rightarrow \mathbb{R}$ defined by

$$h(r) = \frac{\exp(\Lambda R_0^{1-\alpha} r^\alpha)}{r^{\frac{m-1}{2}}} \int_{M \cap B_r} S_1 dM$$

is monotone non-decreasing;

(ii) for $\kappa > 0$ and $\kappa R_0^2 \leq \pi^2$, the function $h : (0, R_0) \rightarrow \mathbb{R}$ defined by

$$h(r) = \frac{\exp(\Lambda R_0^{1-\alpha} r^\alpha)}{(\sin \sqrt{\kappa} r)^{\frac{m-1}{2}}} \int_{M \cap B_r} S_1 dM$$

is monotone non-decreasing.

In particular, if $\overline{M} = \mathbb{R}^{m+1}$ or $\mathbb{S}^{m+1}(\kappa)$ we have, taking $\alpha = 1$ in the Corollary 1.6, the following result.

COROLLARY 1.7 (Monotonicity). *Let M be a proper hypersurface of \mathbb{R}^{m+1} or $\mathbb{S}^{m+1}(\kappa)$, $m \geq 2$, with mean curvature $H > 0$ and scalar curvature $R \geq \kappa$. If there exists $\Lambda \geq 0$ such that*

$$\kappa \leq R \leq \frac{\Lambda}{2} H + \kappa, \quad (1.10)$$

then

(i) for $M^m \subset \mathbb{R}^{m+1}$, the function $\varphi : (0, \infty) \subset \mathbb{R} \rightarrow \mathbb{R}$, defined by

$$\varphi(r) = \frac{e^{\frac{\Lambda}{2} r}}{r^{\frac{m-1}{2}}} \int_{M \cap B_r} H dM$$

is monotone non-decreasing;

(ii) for $M^m \subset \mathbb{S}^{m+1}(\kappa)$, the function $\varphi : \left(0, \frac{\pi}{2\sqrt{\kappa}}\right) \rightarrow \mathbb{R}$ defined by

$$\varphi(r) = \frac{e^{\frac{\Lambda}{2} r}}{(\sin \sqrt{\kappa} r)^{\frac{m-1}{2}}} \int_{M \cap B_r} H dM$$

is monotone non-decreasing.

REMARK 1.5. For the case when $\overline{M} = \mathbb{H}^{m+1}(\kappa)$, results were obtained in [3].

In the next application of the mean value inequalities we study the behaviour of the integral of the mean curvature when we assume L^p bounds for the scalar curvature.

COROLLARY 1.8. *Let \overline{M}^{m+1} , $m \geq 2$, be a Riemannian $(m+1)$ -manifold with sectional curvatures bounded from above by a constant κ . Let M be a proper hypersurface of \overline{M} with $S_1 \geq c$ for some constant $c > 0$ and $S_2 \geq 0$. If there exists $0 < R_0 < i(\overline{M})$, $\Lambda > 0$ and $p > 1$ such that*

$$\left[\int_{M \cap B_{R_0}} \left(\frac{m(\kappa - \kappa_0)}{4} + S_2 \right)^p dM \right]^{1/p} \leq \Lambda, \quad (1.11)$$

then for every $0 < s < t < R_0$, we have

(i) for $\kappa \leq 0$,

$$\begin{aligned} & \left(\frac{1}{s^{\frac{m-1}{2}}} \int_{M \cap B_s} S_1 dM \right)^{1/p} \\ & \leq \left(\frac{1}{t^{\frac{m-1}{2}}} \int_{M \cap B_t} S_1 dM \right)^{1/p} + \frac{2\Lambda}{c^{1-1/p}(m-1-2p)} \int_s^t \frac{1}{r^{\frac{m-1}{2p}}} dr; \end{aligned}$$

(ii) for $\kappa > 0$ and $R_0 \leq \frac{\pi}{2\sqrt{\kappa}}$,

$$\begin{aligned} & \left(\frac{1}{(\sin \sqrt{\kappa} s)^{\frac{m-1}{2}}} \int_{M \cap B_s} S_1 dM \right)^{1/p} \\ & \leq \left(\frac{1}{(\sin \sqrt{\kappa} t)^{\frac{m-1}{2}}} \int_{M \cap B_t} S_1 dM \right)^{1/p} + \frac{\Lambda(m-1)}{c^{1-1/p}(m-1-2p)} \int_s^t \frac{1}{(\sin \sqrt{\kappa} r)^{\frac{m-1}{2p}}} dr. \end{aligned}$$

We conclude the applications of the mean value inequalities with the following monotonicity result for self-shrinkers.

COROLLARY 1.9. *Let M be a proper self-shrinker of \mathbb{R}^{m+1} , $m \geq 2$, with mean curvature $H > 0$ and scalar curvature $0 \leq R \leq \Lambda$ for some constant $\Lambda \geq 0$. Then the function $\varphi : (0, \infty) \rightarrow \mathbb{R}$ defined by*

$$\varphi(r) = \frac{1}{r^{(m-1)(\frac{1}{2}-m\Lambda)}} \int_{M \cap B_r} H dM$$

is monotone non-decreasing. Moreover, if M is complete, non-compact and the scalar curvature satisfies $0 \leq R \leq \Lambda < \frac{1}{2m}$, then $\int_M H dM = \infty$.

This paper is organized in four sections as follows. In the section 2 we present some preliminary results which give basis to establish the argument used in this paper, including Proposition 2.1 and Lemma 2.2. In section 3, we prove the Poincaré type inequalities and, in section 4, we prove the mean value inequalities and its consequences.

Acknowledgements. The authors are grateful to Detang Zhou and Gregório Pacelli Bessa for their suggestions.

2. Preliminary results. Let M be a m -dimensional hypersurface of a Riemannian $(m + 1)$ -manifold \overline{M} , $m \geq 2$. Denote by ∇ and $\overline{\nabla}$ the connections of M and \overline{M} , respectively. Let $\overline{X} : M \rightarrow T\overline{M}$ be a vector field and write $\overline{X} = X^T + X^N$, where $X^T \in TM$ and $X^N \in TM^\perp$. Let $Y, Z \in TM$ be vector fields, and denote by $\langle \cdot, \cdot \rangle$ the metric of \overline{M} . Then

$$\begin{aligned} \langle \overline{\nabla}_Y \overline{X}, Z \rangle &= \langle \overline{\nabla}_Y X^T + \overline{\nabla}_Y X^N, Z \rangle \\ &= \langle \overline{\nabla}_Y X^T, Z \rangle + \langle \overline{\nabla}_Y X^N, Z \rangle \\ &= \langle \overline{\nabla}_Y X^T, Z \rangle - \langle X^N, \overline{\nabla}_Y Z \rangle \\ &= \langle \overline{\nabla}_Y X^T, Z \rangle - \langle X^N, B(Y, Z) \rangle, \end{aligned}$$

where $B(Y, Z) = \overline{\nabla}_Y Z - \nabla_Y Z$ denotes the bilinear form associated with the second fundamental form of M . If η denotes the unit normal vector field of M , then $X^N =$

$\langle \bar{X}, \eta \rangle \eta$. It implies

$$\begin{aligned} \langle \bar{\nabla}_Y \bar{X}, Z \rangle &= \langle \bar{\nabla}_Y X^T, Z \rangle - \langle \bar{X}, \eta \rangle \langle \eta, B(Y, Z) \rangle \\ &= \langle \bar{\nabla}_Y X^T, Z \rangle - \langle \bar{X}, \eta \rangle \langle A(Y), Z \rangle, \end{aligned} \quad (2.1)$$

where $A : TM \rightarrow TM$ is the linear operator defined by

$$\langle A(V), W \rangle = \langle \eta, B(V, W) \rangle, \quad V, W \in TM. \quad (2.2)$$

DEFINITION 2.1. Let $A : TM \rightarrow TM$ be defined by (2.2) the linear operator associated to the second fundamental form of M . The first Newton transformation $P_1 : TM \rightarrow TM$ is defined by

$$P_1 = S_1 I - A,$$

where $S_1 = \text{tr}_M A$, tr_M denotes the trace in M , and $I : TM \rightarrow TM$ is the identity map.

REMARK 2.1. Notice that, since A is self-adjoint, then P_1 is also a self-adjoint linear operator. Denote by k_1, k_2, \dots, k_m the eigenvalues of the linear operator A , also called principal curvatures of M . Since P_1 is a self-adjoint operator, we can consider its eigenvalues $\theta_1, \theta_2, \dots, \theta_m$ given by $\theta_i = S_1 - k_i$, $i = 1, 2, \dots, m$.

REMARK 2.2. If $S_1 > 0$ and $S_2 \geq 0$, then P_1 is positive semidefinite. This fact is known, and can be found in [2, Remark 2.1, p. 552], however, we present a proof here for the sake of completeness. If $S_2 \geq 0$, then $S_1^2 = |A|^2 + 2S_2 \geq k_i^2$, for all $i = 1, 2, \dots, m$. Thus, $0 \leq S_1^2 - k_i^2 = (S_1 - k_i)(S_1 + k_i)$ which implies that all eigenvalues of P_1 are non-negative, provided $S_1 > 0$, i.e., P_1 is positive semidefinite.

The following result is known and we include a proof here for the sake of completeness.

LEMMA 2.1. If $(\text{div}_M P_1)(V) = \text{tr}_M(E \mapsto (\nabla_E P_1)(V))$, where $(\nabla_E P_1)(V) = \nabla_E(P_1(V)) - P_1(\nabla_E V)$, then

$$(\text{div}_M P_1)(V) = \text{Ric}_{\bar{M}}(V, \eta)$$

for every $V \in TM$, where η denotes the unitary normal vector field of M and $\text{Ric}_{\bar{M}}$ denotes the Ricci tensor of \bar{M} . In particular, if \bar{M}^{m+1} has constant sectional curvature or is an Einstein manifold, then $\text{div}_M P_1 = 0$.

Proof. Let $\{e_1, e_2, \dots, e_m\}$ be an orthonormal referential in TM which is geodesic at $p \in M$. By using the Codazzi equation

$$\langle (\nabla_V A)(Y), Z \rangle - \langle (\nabla_Y A)(V), Z \rangle = \langle \bar{R}(Y, V)Z, \eta \rangle$$

for $Y = Z = e_i$ and summing over i from 1 to m , we have

$$\sum_{i=1}^m \langle (\nabla_V A)(e_i), e_i \rangle - \langle (\nabla_{e_i} A)(V), e_i \rangle = \langle \bar{R}(e_i, V)e_i, \eta \rangle,$$

i.e.,

$$\sum_{i=1}^m \langle (\nabla_V A)(e_i), e_i \rangle - (\operatorname{div}_M A)(V) = \operatorname{Ric}_{\overline{M}}(V, \eta).$$

Observing that

$$\sum_{i=1}^m \langle (\nabla_V A)(e_i), e_i \rangle = \sum_{i=1}^m V \langle A(e_i), e_i \rangle = V(S_1) = \operatorname{div}_M(S_1 I)(V),$$

where $I : TM \rightarrow TM$ is the identity operator, we conclude that

$$\operatorname{div}_M(S_1 I - A)(V) = \operatorname{Ric}_{\overline{M}}(V, \eta).$$

In particular, if \overline{M} has constant sectional curvature κ , then $\operatorname{Ric}_{\overline{M}}(V, \eta) = m\kappa \langle V, \eta \rangle = 0$, which implies $\operatorname{div}_M P_1 = 0$. Also, if \overline{M} is an Einstein manifold with Einstein constant λ , we have $\operatorname{Ric}_{\overline{M}}(V, \eta) = \lambda \langle V, \eta \rangle = 0$. Therefore $(\operatorname{div}_M P_1) = 0$. \square

The next result is an important tool in the proof of the Theorem 1.2.

PROPOSITION 2.1. *If $M^m, m \geq 2$, is an hypersurface of a Riemannian $(m + 1)$ -manifold \overline{M}^{m+1} and $\overline{X} : M \rightarrow T\overline{M}$ is a vector field, then*

$$\operatorname{div}_M(P_1(X^T)) = \operatorname{tr}_M \left(E \mapsto P_1 \left((\nabla_E \overline{X})^T \right) \right) + \operatorname{Ric}_{\overline{M}}(X^T, \eta) + 2S_2(\overline{X}, \eta), \tag{2.3}$$

where $\operatorname{Ric}_{\overline{M}}$ denotes the Ricci tensor of \overline{M} and $X^T = \overline{X} - \langle \overline{X}, \eta \rangle \eta$ is the tangent part of \overline{X} .

Proof. Let $\{e_1, e_2, \dots, e_m\}$ be an orthonormal referential in TM . First, since P_1 is self-adjoint, we have

$$\operatorname{tr}_M \left(E \mapsto P_1 \left((\nabla_E \overline{X})^T \right) \right) = \sum_{i=1}^m \langle P_1 \left((\nabla_{e_i} \overline{X})^T \right), e_i \rangle = \sum_{i=1}^m \langle \nabla_{e_i} \overline{X}, P_1(e_i) \rangle. \tag{2.4}$$

By using (2.1), p.705, and the self-adjointness of A , we obtain

$$\begin{aligned} \sum_{i=1}^m \langle \nabla_{e_i} \overline{X}, P_1(e_i) \rangle &= \sum_{i=1}^m \langle \nabla_{e_i} X^T, P_1(e_i) \rangle - \left(\sum_{i=1}^m \langle A(e_i), P_1(e_i) \rangle \right) \langle \overline{X}, \eta \rangle \\ &= \sum_{i=1}^m \langle \nabla_{e_i} X^T, P_1(e_i) \rangle - \left(\sum_{i=1}^m \langle (A \circ P_1)(e_i), e_i \rangle \right) \langle \overline{X}, \eta \rangle \\ &= \sum_{i=1}^m \langle \nabla_{e_i} X^T, P_1(e_i) \rangle - \operatorname{tr}_M(A \circ P_1) \langle \overline{X}, \eta \rangle. \end{aligned}$$

Thus,

$$\sum_{i=1}^m \langle \nabla_{e_i} X^T, P_1(e_i) \rangle = \operatorname{tr}_M \left(E \mapsto P_1 \left((\nabla_E \overline{X})^T \right) \right) + \operatorname{tr}_M(A \circ P_1) \langle \overline{X}, \eta \rangle.$$

On the other hand, the self-adjointness of P_1 implies

$$\begin{aligned}
\sum_{i=1}^m \langle \bar{\nabla}_{e_i} X^T, P_1(e_i) \rangle &= \sum_{i=1}^m \langle \nabla_{e_i} X^T + B(e_i, X^T), P_1(e_i) \rangle = \sum_{i=1}^m \langle \nabla_{e_i} X^T, P_1(e_i) \rangle \\
&= \sum_{i=1}^m \langle P_1(\nabla_{e_i} X^T), e_i \rangle = \sum_{i=1}^m \langle \nabla_{e_i} (P_1(X^T)), e_i \rangle - \sum_{i=1}^m \langle (\nabla_{e_i} P_1)(X^T), e_i \rangle \\
&= \operatorname{div}_M(P_1(X^T)) - \operatorname{tr}_M(E \rightarrow (\nabla_E P_1)(X^T)) \\
&= \operatorname{div}_M(P_1(X^T)) - (\operatorname{div}_M P_1)(X^T).
\end{aligned}$$

Therefore,

$$\operatorname{div}_M(P_1(X^T)) = \operatorname{tr}_M \left(E \mapsto P_1 \left((\bar{\nabla}_E \bar{X})^T \right) \right) + (\operatorname{div}_M P_1)(X^T) + \operatorname{tr}_M(A \circ P_1) \langle \bar{X}, \eta \rangle.$$

The result follows using the Lemma 2.1 and the fact

$$\operatorname{tr}_M(A \circ P_1) = \operatorname{tr}_M(A \circ (S_1 I - A)) = S_1 \operatorname{tr}_M(A) - \operatorname{tr}_M(A^2) = S_1^2 - |A|^2 = 2S_2.$$

□

REMARK 2.3. If the sectional curvatures $K_{\bar{M}}$ of \bar{M}^{m+1} satisfy

$$\kappa_0 \leq K_{\bar{M}} \leq \kappa$$

for real numbers κ_0 e κ , then

$$|\operatorname{Ric}_{\bar{M}}(\bar{V}, \bar{W})| \leq \frac{m(\kappa - \kappa_0)}{2} \quad (2.5)$$

for any orthogonal pair of vectors $\bar{V}, \bar{W} \in T\bar{M}$ such that $|\bar{V}| \leq 1$, $|\bar{W}| \leq 1$. In fact, considering the orthonormal referential $\{\bar{e}_1, \bar{e}_2, \dots, \bar{e}_m, \bar{e}_{m+1}\}$ tangent to \bar{M} , we have

$$\begin{aligned}
\operatorname{Ric}_{\bar{M}}(\bar{V}, \bar{V}) &= \sum_{i=1}^{m+1} \langle \bar{R}(\bar{V}, \bar{e}_i) \bar{V}, \bar{e}_i \rangle = \sum_{i=1}^{m+1} K_{\bar{M}}(\bar{V}, \bar{e}_i) (|\bar{V}|^2 - \langle \bar{V}, \bar{e}_i \rangle^2) \\
&\leq \kappa \sum_{i=1}^{m+1} (|\bar{V}|^2 - \langle \bar{V}, \bar{e}_i \rangle^2) = \kappa [(m+1)|\bar{V}|^2 - |\bar{V}|^2] \\
&= m\kappa |\bar{V}|^2.
\end{aligned}$$

Analogously,

$$\operatorname{Ric}_{\bar{M}}(\bar{V}, \bar{V}) \geq m\kappa_0 |\bar{V}|^2.$$

Now, since $\operatorname{Ric}_{\bar{M}}$ is a symmetric and bilinear form, we have

$$\begin{aligned}
\operatorname{Ric}_{\bar{M}}(\bar{V}, \bar{W}) &= \frac{1}{4} [\operatorname{Ric}_{\bar{M}}(\bar{V} + \bar{W}, \bar{V} + \bar{W}) - \operatorname{Ric}_{\bar{M}}(\bar{V} - \bar{W}, \bar{V} - \bar{W})], \\
&\leq \frac{1}{4} [m\kappa |\bar{V} + \bar{W}|^2 - m\kappa_0 |\bar{V} - \bar{W}|^2] \\
&= \frac{1}{4} [m\kappa (|\bar{V}|^2 + 2\langle \bar{V}, \bar{W} \rangle + |\bar{W}|^2) - m\kappa_0 (|\bar{V}|^2 - 2\langle \bar{V}, \bar{W} \rangle + |\bar{W}|^2)] \\
&= \frac{1}{4} m(\kappa - \kappa_0) (|\bar{V}|^2 + |\bar{W}|^2) \\
&\leq \frac{m(\kappa - \kappa_0)}{2}.
\end{aligned}$$

Analogously,

$$\text{Ric}_{\overline{M}}(\overline{V}, \overline{W}) \geq -\frac{m(\kappa - \kappa_0)}{2}.$$

Therefore,

$$-\frac{m(\kappa - \kappa_0)}{2} \leq \text{Ric}_{\overline{M}}(\overline{V}, \overline{W}) \leq \frac{m(\kappa - \kappa_0)}{2}.$$

This concludes the proof of the estimate (2.5).

In the next lemma we will estimate $\text{tr}_M \left(E \mapsto P_1 \left((\nabla_E \overline{X})^T \right) \right)$ in terms of S_1 , the distance function of \overline{M} and the upper bound κ of the sectional curvatures of \overline{M} .

LEMMA 2.2. *Let \overline{M}^{m+1} , $m \geq 2$, be a Riemannian $(m+1)$ -manifold with sectional curvatures bounded from above by a constant κ , M be a hypersurface of \overline{M}^{m+1} such that $S_1 > 0$ and $S_2 \geq 0$, and $\rho(x) = \rho(p, x)$ be the geodesic distance of \overline{M} starting at a point $x_0 \in \overline{M}$. If $x \in M$ satisfies $\rho(x) < i(\overline{M})$, then*

(i) *for $\kappa \leq 0$ and $\overline{X} = \rho \nabla \rho$,*

$$\text{tr}_M \left(E \mapsto P_1 \left((\nabla_E \overline{X})^T \right) \right) (x) \geq (m - 1)S_1(x);$$

(ii) *for $\kappa > 0$, $\rho(x) < \frac{\pi}{2\sqrt{\kappa}}$ and $\overline{X} = \frac{1}{\sqrt{\kappa}}(\sin \sqrt{\kappa}\rho)\nabla \rho$,*

$$\text{tr}_M \left(E \mapsto P_1 \left((\nabla_E \overline{X})^T \right) \right) (x) \geq (m - 1)S_1(x)(\cos \sqrt{\kappa}\rho(x)).$$

Proof. Let $\gamma : [0, \rho(x)] \rightarrow \overline{M}$ defined by $\gamma(t) = \exp_{x_0}(tu)$, $u \in T_{x_0}\overline{M}$, be the unit speed geodesic such that $\gamma(0) = x_0$ e $\gamma(\rho(x)) = x$. Let $\{e_1(x), e_2(x), \dots, e_m(x)\}$ be an orthonormal basis of $T_x M$ composed by eigenvectors of P_1 in $x \in M$, i.e.,

$$P_1(e_i(x)) = \theta_i(x)e_i(x), \quad i = 1, 2, \dots, m,$$

see Remark 2.1, p.705. Let Y_i , $i = 1, 2, \dots, m$, be the unitary projections of $e_i(x)$ over $\gamma'(\rho(x))^\perp \subset T_x \overline{M}$, namely,

$$Y_i = \frac{e_i(x) - \langle e_i(x), \gamma'(\rho(x)) \rangle \gamma'(\rho(x))}{\|e_i(x) - \langle e_i(x), \gamma'(\rho(x)) \rangle \gamma'(\rho(x))\|}, \quad i = 1, 2, \dots, m.$$

Thus,

$$e_i(x) = b_i Y_i + c_i \gamma'(\rho(x)),$$

where $b_i = \|e_i(x) - \langle e_i(x), \gamma'(\rho(x)) \rangle \gamma'(\rho(x))\|$ and $c_i = \langle e_i(x), \gamma'(\rho(x)) \rangle$ satisfy $b_i^2 + c_i^2 = 1$ and $Y_i \perp \gamma'$ for all $i = 1, 2, \dots, m$. Since the sectional curvatures of \overline{M} satisfy $K_{\overline{M}} \leq \kappa$ and, in the case we assume $\kappa > 0$, we have $\rho(x) < \frac{\pi}{2\sqrt{\kappa}}$, then do not exist

conjugate points to x_0 along γ . Then

$$\begin{aligned} \operatorname{tr}_M \left(E \mapsto P_1 \left((\overline{\nabla}_E \overline{X})^T \right) \right) &= \sum_{i=1}^m \langle \overline{\nabla}_{e_i} \overline{X}, P_1(e_i) \rangle = \sum_{i=1}^m \theta_i \langle \overline{\nabla}_{e_i} \overline{X}, e_i \rangle \\ &= \sum_{i=1}^m \theta_i \langle \overline{\nabla}_{b_i Y_i + c_i \gamma'} \overline{X}, b_i Y_i + c_i \gamma' \rangle \\ &= \sum_{i=1}^m \theta_i b_i^2 \langle \overline{\nabla}_{Y_i} \overline{X}, Y_i \rangle + \sum_{i=1}^m \theta_i c_i^2 \langle \overline{\nabla}_{\gamma'} \overline{X}, \gamma' \rangle \\ &\quad + \sum_{i=1}^m \theta_i b_i c_i [\langle \overline{\nabla}_{Y_i} \overline{X}, \gamma' \rangle + \langle \overline{\nabla}_{\gamma'} \overline{X}, Y_i \rangle]. \end{aligned}$$

In order to unify the proof, let us denote by

$$G(\rho) = \begin{cases} \rho, & \text{if } \kappa \leq 0; \\ \frac{1}{\sqrt{\kappa}} (\sin \sqrt{\kappa} \rho), & \text{if } \kappa > 0. \end{cases}$$

Since $\overline{X}(t) = G(\rho(t)) \overline{\nabla} \rho(t) = G(\rho(t)) \gamma'(t)$ and $\overline{\nabla}_{\gamma'} \gamma' = 0$, we have

$$\begin{aligned} \langle \overline{\nabla}_{\gamma'} \overline{X}, \gamma' \rangle &= \langle \overline{\nabla}_{\gamma'} (G(\rho) \gamma'), \gamma' \rangle = \langle G'(\rho) \langle \overline{\nabla} \rho, \gamma' \rangle \gamma' + G(\rho) \overline{\nabla}_{\gamma'} \gamma', \gamma' \rangle \\ &= G'(\rho) \langle \overline{\nabla} \rho, \gamma' \rangle \langle \gamma', \gamma' \rangle = G'(\rho), \\ \langle \overline{\nabla}_{Y_i} \overline{X}, \gamma' \rangle &= \langle \overline{\nabla}_{Y_i} (G(\rho) \gamma'), \gamma' \rangle = \langle G'(\rho) \langle Y_i, \overline{\nabla} \rho \rangle \gamma' + G(\rho) \overline{\nabla}_{Y_i} \gamma', \gamma' \rangle \\ &= G'(\rho) \langle Y_i, \gamma' \rangle + G(\rho) \langle \overline{\nabla}_{Y_i} \gamma', \gamma' \rangle \\ &= \frac{G(\rho)}{2} Y_i \langle \gamma', \gamma' \rangle = 0, \\ \langle \overline{\nabla}_{\gamma'} \overline{X}, Y_i \rangle &= \langle \overline{\nabla}_{\gamma'} (G(\rho) \gamma'), Y_i \rangle = \langle G'(\rho) \langle \gamma', \overline{\nabla} \rho \rangle \gamma' + G(\rho) \overline{\nabla}_{\gamma'} \gamma', Y_i \rangle = 0. \end{aligned}$$

Thus,

$$\operatorname{tr}_M \left(E \mapsto P_1 \left((\overline{\nabla}_E \overline{X})^T \right) \right) = \sum_{i=1}^m \theta_i b_i^2 \langle \overline{\nabla}_{Y_i} \overline{X}, Y_i \rangle + \sum_{i=1}^m \theta_i c_i^2.$$

On the other hand, is well known that

$$\langle \overline{\nabla}_U \overline{\nabla} \rho, V \rangle = \frac{G'(\rho)}{G(\rho)} [\langle U, V \rangle - \langle U, \overline{\nabla} \rho \rangle \langle V, \overline{\nabla} \rho \rangle],$$

for \mathbb{R}^{m+1} and $\mathbb{S}^{m+1}(\kappa)$. Since

$$\begin{aligned} \langle \overline{\nabla}_{Y_i} \overline{X}, Y_i \rangle &= \langle \overline{\nabla}_{Y_i} (G(\rho) \overline{\nabla} \rho), Y_i \rangle \\ &= \langle G'(\rho) \langle Y_i, \overline{\nabla} \rho \rangle \overline{\nabla} \rho + G(\rho) \overline{\nabla}_{Y_i} \overline{\nabla} \rho, Y_i \rangle \\ &= G(\rho) \langle \overline{\nabla}_{Y_i} \overline{\nabla} \rho, Y_i \rangle, \end{aligned}$$

using Hessian comparison theorem for \overline{M} and the model spaces \mathbb{R}^{m+1} for $\kappa \leq 0$ and $\mathbb{S}^{m+1}(\kappa)$ for $\kappa > 0$, we have

$$\begin{aligned} \langle \overline{\nabla}_{Y_i} \overline{X}, Y_i \rangle &= G(\rho) \langle \overline{\nabla}_{Y_i} \overline{\nabla} \rho, Y_i \rangle \\ &\geq G'(\rho) [|Y_i|^2 - \langle Y_i, \overline{\nabla} \rho \rangle^2] \\ &= G'(\rho). \end{aligned}$$

Therefore,

$$\begin{aligned} \operatorname{tr}_M \left(E \mapsto P_1 \left((\overline{\nabla}_E \overline{X})^T \right) \right) &= \sum_{i=1}^m \theta_i b_i^2 \langle \overline{\nabla}_{Y_i} \overline{X}, Y_i \rangle + \sum_{i=1}^m \theta_i c_i^2 \\ &= G'(\rho) \sum_{i=1}^m \theta_i b_i^2 + \sum_{i=1}^m \theta_i c_i^2 \\ &\geq G'(\rho) \sum_{i=1}^m \theta_i (b_i^2 + c_i^2) = G'(\rho) \sum_{i=1}^m \theta_i \\ &= (m - 1)S_1 G'(\rho). \end{aligned}$$

Since $G'(\rho) = 1$ for $\kappa \leq 0$ and $G'(\rho) = (\cos \sqrt{\kappa} \rho) \leq 1$ for $\kappa > 0$, we have the result. \square

We conclude this section with the following

LEMMA 2.3. *Let M be a hypersurface of a Riemannian $(m + 1)$ -manifold \overline{M}^{m+1} such that $S_1 > 0$ and $S_2 \geq 0$. Let $P_1 = S_1 I - A$, where $A : TM \rightarrow TM$ denotes the linear operator associated to the second fundamental form of M and $I : TM \rightarrow TM$ denotes the identity operator. Then*

$$|\langle P_1(U), V \rangle| \leq 2S_1 |U| |V|$$

for all $U, V \in TM$.

Proof. Let $\theta_i, i = 1, 2, \dots, m$, be the eigenvalues of P_1 . Since $\theta_i = S_1 - k_i$, where k_i are the eigenvalues of the second fundamental form A , we have

$$\begin{aligned} \theta_i = S_1 - k_i &\leq S_1 + |k_i| \\ &\leq S_1 + \sqrt{k_1^2 + k_2^2 + \dots + k_m^2} \\ &= S_1 + |A| = S_1 + \sqrt{S_1^2 - 2S_2} \\ &\leq 2S_1. \end{aligned} \tag{2.6}$$

By using that P_1 is positive semidefinite, the Cauchy-Schwarz inequality, and the estimate (2.6), we obtain

$$\begin{aligned} |\langle P_1(U), V \rangle| &= |\langle \sqrt{P_1}(U), \sqrt{P_1}(V) \rangle| \\ &\leq |\sqrt{P_1}(U)| |\sqrt{P_1}(V)| \\ &= \langle P_1(U), U \rangle^{1/2} \langle P_1(V), V \rangle^{1/2} \\ &\leq (2S_1)^{1/2} |U| (2S_1)^{1/2} |V| \\ &= 2S_1 |U| |V|. \end{aligned} \tag{2.7}$$

\square

3. Proof of the Poincaré inequality.

Proof of Theorem 1.2. Case $\kappa \leq 0$. Initially, since $(\operatorname{diam} \Omega) < 2i(\overline{M})$, we can consider $B_r(x_0), x_0 \in \overline{M}$, the smallest extrinsic ball containing $\overline{\Omega}$, and $\rho(x) = \rho(x_0, x)$ the extrinsic distance from x_0 to $x \in M$. Since $\Omega \subset B_r(x_0)$, then, for all $x \in \Omega$,

$$\rho(x) \leq r = \frac{(\operatorname{diam} \Omega)}{2}. \tag{3.1}$$

Multiplying by f the expression (2.3) in the Proposition 2.1, p. 706, for the vector field $\bar{X} = \rho \bar{\nabla} \rho$ and using item (i) of Lemma 2.2, p. 708, we have

$$f \operatorname{div}_M(P_1(\rho \nabla \rho)) \geq (m-1)fS_1 + f \operatorname{Ric}_{\bar{M}}(\rho \nabla \rho, \eta) + 2S_2 f \langle \rho \bar{\nabla} \rho, \eta \rangle. \quad (3.2)$$

This implies

$$\begin{aligned} \operatorname{div}_M(fP_1(\rho \nabla \rho)) &= f \operatorname{div}_M(P_1(\rho \nabla \rho)) + \langle \nabla f, P_1(\rho \nabla \rho) \rangle \\ &\geq (m-1)fS_1 + f \operatorname{Ric}_{\bar{M}}(\rho \nabla \rho, \eta) + 2S_2 f \langle \rho \bar{\nabla} \rho, \eta \rangle + \langle \nabla f, P_1(\rho \nabla \rho) \rangle. \end{aligned}$$

Integrating expression above over Ω , using divergence theorem, we have

$$\begin{aligned} 0 &\geq (m-1) \int_{\Omega} f S_1 dM + \int_{\Omega} f \operatorname{Ric}_{\bar{M}}(\rho \nabla \rho, \eta) dM \\ &\quad + 2 \int_{\Omega} S_2 f \langle \rho \bar{\nabla} \rho, \eta \rangle dM + \int_{\Omega} \langle \nabla f, P_1(\rho \nabla \rho) \rangle dM, \end{aligned}$$

for every function f compactly supported on Ω , i.e.,

$$\begin{aligned} \int_{\Omega} f S_1 dM &\leq \frac{1}{m-1} \left[\int_{\Omega} \langle \nabla f, P_1(-\rho \nabla \rho) \rangle dM + \int_{\Omega} f \operatorname{Ric}_{\bar{M}}(-\rho \nabla \rho, \eta) dM \right. \\ &\quad \left. + 2 \int_{\Omega} S_2 f \langle -\rho \bar{\nabla} \rho, \eta \rangle dM \right]. \end{aligned} \quad (3.3)$$

By using Lemma 2.3 for $U = -\rho \nabla \rho$ and $V = \nabla f$, we have

$$|\langle \nabla f, P_1(-\rho \nabla \rho) \rangle| \leq 2S_1 \rho |\nabla f|,$$

and the estimate (2.5), p. 707, gives

$$\operatorname{Ric}_{\bar{M}}(-\nabla \rho, \eta) \leq \frac{m(\kappa - \kappa_0)}{2}.$$

Replacing these inequalities in (3.3) we obtain

$$\int_{\Omega} f S_1 dM \leq \frac{2}{m-1} \int_{\Omega} \rho \left[|\nabla f| S_1 + \left(\frac{m(\kappa - \kappa_0)}{4} + S_2 \right) f \right] dM.$$

Therefore, using (3.1),

$$\int_{\Omega} f S_1 dM \leq \frac{1}{m-1} (\operatorname{diam} \Omega) \int_{\Omega} \left[|\nabla f| S_1 + \left(\frac{m(\kappa - \kappa_0)}{4} + S_2 \right) f \right] dM.$$

Case $\kappa > 0$. Again, since $(\operatorname{diam} \Omega) < 2i(\bar{M})$ and $(\operatorname{diam} \Omega) < \frac{\pi}{\sqrt{\kappa}}$, we can consider $B_r(x_0)$, $x_0 \in \bar{M}^{m+1}$, the smallest extrinsic ball containing $\bar{\Omega}$, and $\rho(x) = \rho(x_0, x)$ the extrinsic distance from x_0 to $x \in M$. Since $\Omega \subset B_r(x_0)$, then

$$\rho(x) \leq r = \frac{(\operatorname{diam} \Omega)}{2}, \quad (3.4)$$

for all $x \in \Omega$. By using the Proposition 2.1 for $\bar{X} = \frac{1}{\sqrt{\kappa}} (\sin \sqrt{\kappa} \rho) \bar{\nabla} \rho$ and using the Lemma 2.2, item (ii), we have

$$\begin{aligned} f \operatorname{div}_M \left(P_1 \left(\frac{(\sin \sqrt{\kappa} \rho)}{\sqrt{\kappa}} \nabla \rho \right) \right) &\geq (m-1)fS_1(\cos \sqrt{\kappa} \rho) + \frac{(\sin \sqrt{\kappa} \rho)}{\sqrt{\kappa}} f \operatorname{Ric}_{\bar{M}}(\nabla \rho, \eta) \\ &\quad + 2S_2 f \frac{(\sin \sqrt{\kappa} \rho)}{\sqrt{\kappa}} \langle \bar{\nabla} \rho, \eta \rangle. \end{aligned}$$

This implies that

$$\begin{aligned} \operatorname{div}_M(fP_1((\sin \sqrt{\kappa}\rho)\nabla\rho)) &= f \operatorname{div}_M(P_1((\sin \sqrt{\kappa}\rho)\nabla\rho)) + (\sin \sqrt{\kappa}\rho)\langle \nabla f, P_1(\nabla\rho) \rangle \\ &\geq (m-1)fS_1\sqrt{\kappa}(\cos \sqrt{\kappa}\rho) + (\sin \sqrt{\kappa}\rho)f \operatorname{Ric}_{\overline{M}}(\nabla\rho, \eta) \\ &\quad + 2S_2f(\sin \sqrt{\kappa}\rho)\langle \overline{\nabla}\rho, \eta \rangle + (\sin \sqrt{\kappa}\rho)\langle \nabla f, P_1(\nabla\rho) \rangle. \end{aligned}$$

Integrating the expression above over Ω and applying divergence theorem, we obtain

$$\begin{aligned} 0 &\geq (m-1)\sqrt{\kappa} \int_{\Omega} fS_1(\cos \sqrt{\kappa}\rho)dM + \int_{\Omega} (\sin \sqrt{\kappa}\rho)f \operatorname{Ric}_{\overline{M}}(\nabla\rho, \eta)dM \\ &\quad + \int_{\Omega} 2S_2f(\sin \sqrt{\kappa}\rho)\langle \overline{\nabla}\rho, \eta \rangle dM + \int_{\Omega} (\sin \sqrt{\kappa}\rho)\langle \nabla f, P_1(\nabla\rho) \rangle dM, \end{aligned} \tag{3.5}$$

since f is compactly supported in Ω . By using the Lemma 2.3, p. 710, for $U = -\nabla\rho$ and $V = \nabla f$, we have

$$|\langle \nabla f, P_1(-\nabla\rho) \rangle| \leq 2S_1|\nabla f|.$$

Replacing this estimate in (3.5), and using the estimate (2.5), p. 707, we obtain

$$\begin{aligned} \int_{\Omega} fS_1(\cos \sqrt{\kappa}\rho)dM &\leq \frac{1}{(m-1)\sqrt{\kappa}} \int_{\Omega} (\sin \sqrt{\kappa}\rho) [\langle \nabla f, P_1(\nabla\rho) \rangle + f \operatorname{Ric}_{\overline{M}}(\nabla\rho, \eta) + 2S_2f] dM \\ &\leq \frac{2}{(m-1)\sqrt{\kappa}} \int_{\Omega} (\sin \sqrt{\kappa}\rho) \left[|\nabla f|S_1 + \left(\frac{m(\kappa - \kappa_0)}{4} + S_2 \right) f \right] dM. \end{aligned}$$

Since $\rho(x) \leq r < \frac{\pi}{2\sqrt{\kappa}}$ for all $x \in \Omega$, $(\cos \sqrt{\kappa}\rho)$ is a decreasing function and $(\sin \sqrt{\kappa}\rho)$ is an increasing function for $\rho \in \left(0, \frac{\pi}{2\sqrt{\kappa}}\right)$, we have

$$\begin{aligned} (\cos \sqrt{\kappa}r) \int_{\Omega} fS_1dM &\leq \int_{\Omega} fS_1(\cos \sqrt{\kappa}\rho)dM \\ &\leq \frac{2}{(m-1)\sqrt{\kappa}} \int_{\Omega} (\sin \sqrt{\kappa}\rho) \left[|\nabla f|S_1 + \left(\frac{m(\kappa - \kappa_0)}{4} + S_2 \right) f \right] dM \\ &\leq \frac{2}{(m-1)\sqrt{\kappa}} (\sin \sqrt{\kappa}r) \int_{\Omega} \left[|\nabla f|S_1 + \left(\frac{m(\kappa - \kappa_0)}{4} + S_2 \right) f \right] dM. \end{aligned}$$

i.e.,

$$\int_{\Omega} fS_1dM \leq \frac{2(\tan \sqrt{\kappa}r)}{\sqrt{\kappa}(m-1)} \int_{\Omega} \left[|\nabla f|S_1 + \left(\frac{m(\kappa - \kappa_0)}{4} + S_2 \right) f \right] dM.$$

Therefore, using (3.4),

$$\int_{\Omega} fS_1dM \leq \frac{2 \tan \left(\frac{\sqrt{\kappa}}{2}(\operatorname{diam} \Omega) \right)}{\sqrt{\kappa}(m-1)} \int_{\Omega} \left[|\nabla f|S_1 + \left(\frac{m(\kappa - \kappa_0)}{4} + S_2 \right) f \right] dM,$$

for $(\operatorname{diam} \Omega) < \frac{\pi}{\sqrt{\kappa}}$. \square

4. Proof of the mean value inequalities and its applications. From now on, we will let $B_r = B_r(x_0)$ be the extrinsic open ball with center $x_0 \in \overline{M}^{m+1}$ and radius r . If $\partial M \neq \emptyset$, assume in addition that $B_r(x_0) \cap \partial M = \emptyset$.

Proof of Theorem 1.4. Let \overline{X} be a vector field on the ambient space. Since

$$\operatorname{div}_M(P_1(fX^T)) = \langle \nabla f, P_1(X^T) \rangle + f \operatorname{div}_M(P_1(X^T)),$$

using the Proposition 2.1, p. 706, we have

$$\begin{aligned} & \operatorname{div}_M(P_1(fX^T)) \\ &= \langle \nabla f, P_1(X^T) \rangle + f \operatorname{tr}_M(E \mapsto P_1((\overline{\nabla}_E \overline{X})^T)) + f \operatorname{Ric}_{\overline{M}}(X^T, \eta) + 2S_2 f \langle \overline{X}, \eta \rangle. \end{aligned}$$

By using integration, the divergence theorem, Lemma 2.3, and the co-area formula, we have

$$\begin{aligned} \int_{M \cap B_r} \operatorname{div}_M(P_1(fX^T)) dM &= \int_{\partial(M \cap B_r)} \langle P_1(fX^T), \nu \rangle dS_M \\ &\leq 2 \int_{\partial(M \cap B_r)} f S_1 |X^T| dS_M \\ &\leq 2 \sup_{\partial(M \cap B_r)} |X^T| \int_{\partial(M \cap B_r)} f S_1 |\nabla \rho|^{-1} dS_M \\ &= 2 \sup_{\partial(M \cap B_r)} |X^T| \frac{d}{dr} \left(\int_{M \cap B_r} f S_1 dM \right), \end{aligned}$$

where ν is the outer conormal vector field of $\partial(M \cap B_r)$ and dS_M is the volume element of $\partial(M \cap B_r)$. This implies

$$\begin{aligned} & 2 \sup_{\partial(M \cap B_r)} |X^T| \frac{d}{dr} \left(\int_{M \cap B_r} f S_1 dM \right) \\ &\geq \int_{M \cap B_r} \langle \nabla f, P_1(X^T) \rangle dM \\ &\quad + \int_{M \cap B_r} f \operatorname{tr}_M(E \mapsto P_1((\overline{\nabla}_E \overline{X})^T)) dM + \int_{M \cap B_r} f \operatorname{Ric}_{\overline{M}}(X^T, \eta) dM \\ &\quad + 2 \int_{M \cap B_r} S_2 f \langle \overline{X}, \eta \rangle dM. \end{aligned}$$

Case $\kappa \leq 0$. Taking $\overline{X} = \rho \overline{\nabla} \rho$, we have $\sup_{\partial(M \cap B_r)} |X^T| = r$ and, using Lemma 2.2, p. 708, we have

$$\begin{aligned} 2r \frac{d}{dr} \left(\int_{M \cap B_r} f S_1 dM \right) &\geq \int_{M \cap B_r} \langle \nabla f, P_1(\rho \nabla \rho) \rangle dM + (m-1) \int_{M \cap B_r} f S_1 dM \\ &\quad + \int_{M \cap B_r} f \operatorname{Ric}_{\overline{M}}(\rho \nabla \rho, \eta) dM + 2 \int_{M \cap B_r} S_2 f \langle \rho \overline{\nabla} \rho, \eta \rangle dM, \end{aligned}$$

i.e.,

$$\begin{aligned} & r \frac{d}{dr} \left(\int_{M \cap B_r} f S_1 dM \right) - \frac{m-1}{2} \int_{M \cap B_r} f S_1 dM \\ &\geq \frac{1}{2} \int_{M \cap B_r} [\langle \rho \overline{\nabla} \rho, P_1(\nabla f) \rangle + 2S_2 f \eta] + f \operatorname{Ric}_{\overline{M}}(\rho \nabla \rho, \eta) dM. \end{aligned}$$

Since

$$r \frac{d}{dr} \left(\int_{M \cap B_r} f S_1 dM \right) - \frac{m-1}{2} \int_{M \cap B_r} f S_1 dM = r^{\frac{m+1}{2}} \frac{d}{dr} \left(\frac{1}{r^{\frac{m-1}{2}}} \int_{M \cap B_r} f S_1 dM \right),$$

we have, dividing by $r^{\frac{m+1}{2}}$,

$$\begin{aligned} & \frac{d}{dr} \left(\frac{1}{r^{\frac{m-1}{2}}} \int_{M \cap B_r} f S_1 dM \right) \\ & \geq \frac{1}{2r^{\frac{m+1}{2}}} \int_{M \cap B_r} [\langle \rho \bar{\nabla} \rho, P_1(\nabla f) + 2S_2 f \eta \rangle + f \operatorname{Ric}_{\bar{M}}(\rho \nabla \rho, \eta)] dM. \end{aligned} \tag{4.1}$$

Integrating the inequality (4.1) from s to t , we have the result for the case $\kappa \leq 0$.

Case $\kappa > 0$. Consider $\bar{X} = \frac{1}{\sqrt{\kappa}}(\sin \sqrt{\kappa} \rho) \bar{\nabla} \rho$. Since we are assuming $\rho < \frac{\pi}{2\sqrt{\kappa}}$, we have

$$\sup_{\partial(M \cap B_r)} |X^T| = \frac{1}{\sqrt{\kappa}}(\sin \sqrt{\kappa} r)$$

and, using Lemma 2.2, we obtain

$$\begin{aligned} & \frac{2}{\sqrt{\kappa}}(\sin \sqrt{\kappa} r) \frac{d}{dr} \left(\int_{M \cap B_r} f S_1 dM \right) \\ & \geq \int_{M \cap B_r} \frac{1}{\sqrt{\kappa}}(\sin \sqrt{\kappa} \rho) \langle \nabla \rho, P_1(\nabla f) \rangle dM \\ & \quad + (m-1) \int_{M \cap B_r} (\cos \sqrt{\kappa} \rho) f S_1 dM + \int_{M \cap B_r} \frac{(\sin \sqrt{\kappa} \rho)}{\sqrt{\kappa}} f \operatorname{Ric}_{\bar{M}}(\nabla \rho, \eta) dM \\ & \quad + 2 \int_{M \cap B_r} \frac{(\sin \sqrt{\kappa} \rho)}{\sqrt{\kappa}} S_2 f \langle \bar{\nabla} \rho, \eta \rangle dM. \end{aligned}$$

Since $(\cos \sqrt{\kappa} \rho)$ is a decreasing function for $\rho \leq \frac{\pi}{2\sqrt{\kappa}}$, we have

$$\int_{M \cap B_r} (\cos \sqrt{\kappa} \rho) f S_1 dM \geq (\cos \sqrt{\kappa} r) \int_{M \cap B_r} f S_1 dM,$$

and thus, dividing by $\frac{2}{\sqrt{\kappa}}(\sin \sqrt{\kappa} r)$ we obtain

$$\begin{aligned} & \frac{d}{dr} \left(\int_{M \cap B_r} f S_1 dM \right) - \frac{m-1}{2} \sqrt{\kappa} (\cot \sqrt{\kappa} r) \int_{M \cap B_r} f S_1 dM \\ & \geq \frac{1}{2(\sin \sqrt{\kappa} r)} \int_{M \cap B_r} (\sin \sqrt{\kappa} \rho) [\langle \bar{\nabla} \rho, P_1(\nabla f) + 2S_2 f \eta \rangle + f \operatorname{Ric}_{\bar{M}}(\nabla \rho, \eta)] dM. \end{aligned}$$

Since

$$\begin{aligned} & \frac{d}{dr} \left(\int_{M \cap B_r} f S_1 dM \right) - \frac{m-1}{2} \sqrt{\kappa} (\cot \sqrt{\kappa} r) \int_{M \cap B_r} f S_1 dM \\ & = (\sin \sqrt{\kappa} r)^{\frac{m-1}{2}} \frac{d}{dr} \left(\frac{1}{(\sin \sqrt{\kappa} r)^{\frac{m-1}{2}}} \int_{M \cap B_r} f S_1 dM \right), \end{aligned}$$

we have

$$\begin{aligned} & \frac{d}{dr} \left(\frac{1}{(\sin \sqrt{\kappa} r)^{\frac{m-1}{2}}} \int_{M \cap B_r} f S_1 dM \right) \\ & \geq \frac{1}{2(\sin \sqrt{\kappa} r)^{\frac{m+1}{2}}} \int_{M \cap B_r} (\sin \sqrt{\kappa} \rho) [\langle \bar{\nabla} \rho, P_1(\nabla f) + 2S_2 f \eta \rangle + f \operatorname{Ric}_{\bar{M}}(\nabla \rho, \eta)] dM. \end{aligned} \tag{4.2}$$

Integrating the inequality (4.2) from s to t , we have the result for the case $\kappa > 0$. \square

REMARK 4.1. If $A \geq 0$, then we can estimate the eigenvalues of P_1 by $\theta_i = S_1 - k_i \leq S_1$ in the place of $\theta_i \leq 2S_1$ in the proof of the Lemma 2.3, p. 710. In this case, the exponents of the mean value inequalities become $(m-1)$ in the place of $\frac{m-1}{2}$ and the mean value inequalities become:

(i) for $\kappa \leq 0$,

$$\begin{aligned} & \frac{1}{t^{m-1}} \int_{M \cap B_t} f S_1 dM - \frac{1}{s^{m-1}} \int_{M \cap B_s} f S_1 dM \\ & \geq \int_s^t \frac{1}{r^m} \int_{M \cap B_r} [\langle \rho \bar{\nabla} \rho, P_1(\nabla f) + 2S_2 f \eta \rangle + f \operatorname{Ric}_{\bar{M}}(\rho \nabla \rho, \eta)] dM dr; \end{aligned}$$

(ii) for $\kappa > 0$ and $t < \frac{\pi}{2\sqrt{\kappa}}$,

$$\begin{aligned} & \frac{1}{(\sin \sqrt{\kappa} t)^{m-1}} \int_{M \cap B_t} f S_1 dM - \frac{1}{(\sin \sqrt{\kappa} s)^{m-1}} \int_{M \cap B_s} f S_1 dM \\ & \geq \int_s^t \frac{1}{(\sin \sqrt{\kappa} r)^m} \int_{M \cap B_r} (\sin \sqrt{\kappa} \rho) [\langle \bar{\nabla} \rho, P_1(\nabla f) + 2S_2 f \eta \rangle + f \operatorname{Ric}_{\bar{M}}(\nabla \rho, \eta)] dM dr. \end{aligned}$$

Now will prove the corollaries stated in the introduction of this paper.

Proof of Corollary 1.6. Let us prove the case $\kappa > 0$. The case $\kappa \leq 0$ is entirely analogous. Applying the mean value inequality of Theorem 1.4, p. 702, item (ii), for $f \equiv 1$, we have

$$\begin{aligned} & \frac{1}{(\sin \sqrt{\kappa} t)^{\frac{m-1}{2}}} \int_{M \cap B_t} S_1 dM - \frac{1}{(\sin \sqrt{\kappa} s)^{\frac{m-1}{2}}} \int_{M \cap B_s} S_1 dM \\ & \geq \int_s^t \frac{1}{(\sin \sqrt{\kappa} r)^{\frac{m+1}{2}}} \int_{M \cap B_r} (\sin \sqrt{\kappa} \rho) [2S_2 \langle \bar{\nabla} \rho, \eta \rangle + \operatorname{Ric}_{\bar{M}}(\nabla \rho, \eta)] dM dr. \end{aligned}$$

Since

$$\begin{aligned} & \int_{M \cap B_r} (\sin \sqrt{\kappa} \rho) [2S_2 \langle \bar{\nabla} \rho, \eta \rangle + \operatorname{Ric}_{\bar{M}}(\nabla \rho, \eta)] dM \\ & \geq -2(\sin \sqrt{\kappa} r) \int_{M \cap B_r} \left(\frac{m(\kappa - \kappa_0)}{4} + S_2 \right) dM, \end{aligned}$$

and using the hypothesis (1.9), p. 703, we have, for $0 < s < t < R_0$,

$$\begin{aligned} & \frac{1}{(\sin \sqrt{\kappa} t)^{\frac{m-1}{2}}} \int_{M \cap B_t} S_1 dM - \frac{1}{(\sin \sqrt{\kappa} s)^{\frac{m-1}{2}}} \int_{M \cap B_s} S_1 dM \\ & \geq - \int_s^t (\sin \sqrt{\kappa} r)^{-\frac{m-1}{2}} \int_{M \cap B_r} \left(\frac{m(\kappa - \kappa_0)}{4} + S_2 \right) dM dr \\ & \geq -\alpha \Lambda \int_s^t (\sin \sqrt{\kappa} r)^{-\frac{m-1}{2}} \left(\frac{r}{R_0} \right)^{\alpha-1} \int_{M \cap B_r} S_1 dM dr. \end{aligned}$$

Letting $g(r) = \frac{1}{(\sin \sqrt{\kappa} r)^{\frac{m-1}{2}}} \int_{M \cap B_r} S_1 dM$ and dividing by $t - s$, inequality above

becomes

$$\begin{aligned} \frac{g(t) - g(s)}{t - s} &\geq -\frac{1}{t - s} \int_s^t \alpha \Lambda \left(\frac{r}{R_0} \right)^{\alpha-1} g(r) dr \\ &= -\frac{1}{t - s} \left[\int_\varepsilon^t \alpha \Lambda \left(\frac{r}{R_0} \right)^{\alpha-1} g(r) dr - \int_\varepsilon^s \alpha \Lambda \left(\frac{r}{R_0} \right)^{\alpha-1} g(r) dr \right], \end{aligned}$$

for every $\varepsilon > 0$ sufficiently small.

Since $r \mapsto \int_{M \cap B_r} S_1 dM$ is a monotone non-decreasing function, a classical theorem of Lebesgue guarantee that this function is differentiable almost everywhere with respect to Lebesgue measure of \mathbb{R} . Consequently, the same holds for $g(r)$. Considering the points s such that g is differentiable and taking $t \rightarrow s$, the left hand side goes to $g'(s)$ and the right hand side goes to $\alpha \Lambda \left(\frac{s}{R_0} \right)^{\alpha-1} g(s)$. Thus, g satisfies

$$g'(r) + \alpha \Lambda \left(\frac{r}{R_0} \right)^{\alpha-1} g(r) \geq 0.$$

Since

$$\frac{d}{dr} \left(\exp(\Lambda R_0^{1-\alpha} r^\alpha) g(r) \right) = \exp(\Lambda R_0^{1-\alpha} r^\alpha) \left(\alpha \Lambda \left(\frac{r}{R_0} \right)^{\alpha-1} g(r) + g'(r) \right) \geq 0,$$

we conclude that $h(r) = \exp(\Lambda R_0^{1-\alpha} r^\alpha) g(r)$ is monotone non-decreasing for every $r \in (0, R_0)$. \square

Proof of Corollary 1.8. Again, we prove the case $\kappa > 0$. The case $\kappa \leq 0$ is entirely analogous. Applying inequality (4.2) to $f \equiv 1$ and using that $(\sin \sqrt{\kappa} \rho)$ is a increasing function, we have

$$\begin{aligned} &\frac{d}{dr} \left(\frac{1}{(\sin \sqrt{\kappa} r)^{\frac{m-1}{2}}} \int_{M \cap B_r} S_1 dM \right) \\ &\geq \frac{1}{2(\sin \sqrt{\kappa} r)^{\frac{m+1}{2}}} \int_{M \cap B_r} (\sin \sqrt{\kappa} \rho) [2S_2 \langle \nabla \rho, \eta \rangle + \text{Ric}_{\overline{M}}(\nabla \rho, \eta)] dM. \\ &\geq -\frac{1}{(\sin \sqrt{\kappa} r)^{\frac{m-1}{2}}} \int_{M \cap B_r} \left(\frac{m(\kappa - \kappa_0)}{4} + S_2 \right) dM. \end{aligned}$$

By using Hölder inequality and the hypothesis, we obtain

$$\begin{aligned} &\int_{M \cap B_r} \left(\frac{m(\kappa - \kappa_0)}{4} + S_2 \right) dM \\ &\leq \left(\int_{M \cap B_r} \left(\frac{m(\kappa - \kappa_0)}{4} + S_2 \right)^p dM \right)^{\frac{1}{p}} \left(\int_{M \cap B_r} dM \right)^{1-\frac{1}{p}} \\ &\leq \frac{1}{c^{1-\frac{1}{p}}} \left(\int_{M \cap B_{R_0}} \left(\frac{m(\kappa - \kappa_0)}{4} + S_2 \right)^p dM \right)^{\frac{1}{p}} \left(\int_{M \cap B_r} S_1 dM \right)^{1-\frac{1}{p}} \\ &\leq \frac{\Lambda}{c^{1-\frac{1}{p}}} \left(\int_{M \cap B_r} S_1 dM \right)^{1-\frac{1}{p}} dM. \end{aligned}$$

This implies

$$\frac{d}{dr} \left(\frac{1}{(\sin \sqrt{\kappa r})^{\frac{m-1}{2}}} \int_{M \cap B_r} S_1 dM \right) \geq -\frac{\Lambda}{c^{1-1/p}} \frac{1}{(\sin \sqrt{\kappa r})^{\frac{m-1}{2}}} \left(\int_{M \cap B_r} S_1 dM \right)^{1-\frac{1}{p}}.$$

Thus,

$$\begin{aligned} \frac{d}{dr} \left(\left(\frac{1}{(\sin \sqrt{\kappa r})^{\frac{m-1}{2}}} \int_{M \cap B_r} S_1 dM \right)^{\frac{1}{p}} \right) &= \frac{1}{p} \left(\frac{1}{(\sin \sqrt{\kappa r})^{\frac{m-1}{2}}} \int_{M \cap B_r} S_1 dM \right)^{\frac{1}{p}-1} \\ &\quad \times \frac{d}{dr} \left(\frac{1}{(\sin \sqrt{\kappa r})^{\frac{m-1}{2}}} \int_{M \cap B_r} S_1 dM \right) \\ &\geq -\frac{1}{p} \left(\frac{1}{(\sin \sqrt{\kappa r})^{\frac{m-1}{2}}} \int_{M \cap B_r} S_1 dM \right)^{\frac{1}{p}-1} \\ &\quad \times \frac{\Lambda}{c^{1-1/p}} \frac{1}{(\sin \sqrt{\kappa r})^{\frac{m-1}{2}}} \left(\int_{M \cap B_r} S_1 dM \right)^{1-\frac{1}{p}} \\ &= -\frac{\Lambda}{pc^{1-1/p}} \frac{1}{(\sin \sqrt{\kappa r})^{\frac{m-1}{2p}}}. \end{aligned}$$

Integrating inequality above from s to t , we have

$$\begin{aligned} &\left(\frac{1}{(\sin \sqrt{\kappa t})^{\frac{m-1}{2}}} \int_{M \cap B_t} S_1 dM \right)^{\frac{1}{p}} - \left(\frac{1}{(\sin \sqrt{\kappa s})^{\frac{m-1}{2}}} \int_{M \cap B_s} S_1 dM \right)^{\frac{1}{p}} \\ &\geq -\frac{\Lambda}{pc^{1-1/p}} \int_s^t \frac{1}{(\sin \sqrt{\kappa r})^{\frac{m-1}{2p}}} dr. \end{aligned}$$

□

We conclude this paper proving Corollary 1.9, p. 704.

Proof of Corollary 1.9. Taking $f \equiv 1$ in the inequality (4.1), p. 714, for $\overline{M} = \mathbb{R}^{m+1}$, using that M is a self-shrinker and the hypothesis $0 \leq R \leq \Lambda$, we have

$$\begin{aligned} \frac{d}{dr} \left(\frac{1}{r^{\frac{m-1}{2}}} \int_{M \cap B_r} HdM \right) &\geq \frac{m-1}{r^{\frac{m+1}{2}}} \int_{M \cap B_r} R \left(\frac{1}{2} \langle \rho \overline{\nabla} \rho, \eta \rangle \right) dM \\ &= -\frac{m(m-1)}{r^{\frac{m+1}{2}}} \int_{M \cap B_r} RHdM \\ &\geq -\frac{\Lambda m(m-1)}{r^{\frac{m+1}{2}}} \int_{M \cap B_r} HdM. \end{aligned} \tag{4.3}$$

Denoting by $f(r) = \frac{1}{r^{\frac{m-1}{2}}} \int_{M \cap B_r} HdM$, the inequality (4.3) becomes

$$f'(r) \geq -\frac{\Lambda m(m-1)}{r} f(r),$$

which is equivalent to

$$\frac{d}{dr} \left(\ln \left(r^{\Lambda m(m-1)} f(r) \right) \right) \geq 0.$$

This implies that $\ln(r^{\Lambda m(m-1)}f(r))$. Therefore $r^{\Lambda m(m-1)}f(r)$ is monotone non-decreasing, i.e.,

$$\varphi(r) = \frac{1}{r^{(m-1)(\frac{1}{2}-\Lambda m)}} \int_{M \cap B_r} HdM$$

is monotone non-decreasing. The monotonicity of the function φ implies

$$\int_{M \cap B_r} HdM \geq r^{(m-1)(\frac{1}{2}-\Lambda m)} \frac{1}{r_0^{(m-1)(\frac{1}{2}-\Lambda m)}} \int_{M \cap B_{r_0}} HdM$$

for $r \geq r_0$. Therefore, in the case that $0 \leq R \leq \Lambda < \frac{1}{2m}$, taking $r \rightarrow \infty$ we obtain $\int_M HdM = \infty$. \square

REFERENCES

- [1] G. ACOSTA AND R. G. DURÁN, *An optimal Poincaré inequality in L^1 for convex domains*, Proc. Amer. Math. Soc., 132:1 (2004), pp. 195–202 (electronic).
- [2] H. ALENCAR, M. DO CARMO, AND W. SANTOS, *A gap theorem for hypersurfaces of the sphere with constant scalar curvature one*, Comment. Math. Helv., 77:3 (2002), pp. 549–562.
- [3] H. ALENCAR AND G. S. NETO, *Monotonicity formula for complete hypersurfaces in the hyperbolic space and applications*, Arkiv för Matematik, to appear in print (2015).
- [4] C. S. BARROSO, L. L. DE LIMA, AND W. SANTOS, *Monotonicity inequalities for the r -area and a degeneracy theorem for r -minimal graphs*, J. Geom. Anal., 14:4 (2004), pp. 557–566.
- [5] Y. D. BURAGO AND V. A. ZALGALLER, *Geometric inequalities, Translated from the Russian by A. B. Sosinskii*, 2nd ed., Grundlehren der Mathematischen Wissenschaften, Vol. 285, Springer Series in Soviet Mathematics. Springer-Verlag, Berlin, 1988.
- [6] H.-D. CAO AND D. ZHOU, *On complete gradient shrinking Ricci solitons*, J. Differential Geom., 85:2 (2010), pp. 175–185.
- [7] X. CHENG AND D. ZHOU, *Manifolds with weighted Poincaré inequality and uniqueness of minimal hypersurfaces*, Comm. Anal. Geom., 17:1 (2009), pp. 139–154.
- [8] T. H. COLDING AND W. P. MINICOZZI II, *A course in minimal surfaces*, Graduate Studies in Mathematics, Vol. 121, American Mathematical Society, Providence, RI, 2011.
- [9] T. H. COLDING, *New monotonicity formulas for Ricci curvature and applications. I*, Acta Math., 209:2 (2012), pp. 229–263.
- [10] T. H. COLDING AND W. P. MINICOZZI II, *Ricci curvature and monotonicity for harmonic functions*, Calc. Var. Partial Differential Equations, 49:3-4 (2014), pp. 1045–1059.
- [11] ———, *Generic mean curvature flow I: generic singularities*, Ann. of Math. (2), 175:2 (2012), pp. 755–833.
- [12] ———, *Smooth compactness of self-shrinkers*, Comment. Math. Helv., 87:2 (2012), pp. 463–475.
- [13] M. P. DO CARMO, *Riemannian geometry*, Mathematics: Theory & Applications, Birkhäuser Boston, Inc., Boston, MA, 1992. Translated from the second Portuguese edition by Francis Flaherty.
- [14] K. ECKER, *A local monotonicity formula for mean curvature flow*, Ann. of Math. (2), 154:2 (2001), pp. 503–525.
- [15] ———, *Local monotonicity formulas for some nonlinear diffusion equations*, Calc. Var. Partial Differential Equations, 23:1 (2005), pp. 67–81.
- [16] L. C. EVANS, *Partial differential equations*, 2nd ed., Graduate Studies in Mathematics, Vol. 19, American Mathematical Society, Providence, RI, 2010.
- [17] P. GUAN AND J. LI, *A mean curvature type flow in space forms*, International Mathematics Research Notices, 2015:13 (2014), pp. 4716–4740.
- [18] M. GRÜTER, *The monotonicity formula in geometric measure theory, and an application to a partially free boundary problem*, Partial differential equations and calculus of variations, Lecture Notes in Math., Vol. 1357, Springer, Berlin, 1988 pp. 238–255.
- [19] D. HOFFMAN AND J. SPRUCK, *Sobolev and isoperimetric inequalities for Riemannian submanifolds*, Comm. Pure Appl. Math., 27 (1974), pp. 715–727.
- [20] I. KOMBE AND M. ÖZAYDIN, *Hardy-Poincaré, Rellich and uncertainty principle inequalities on Riemannian manifolds*, Trans. Amer. Math. Soc., 365:10 (2013), pp. 5035–5050.

- [21] K.-H. LAM, *Results on a weighted Poincaré inequality of complete manifolds*, Trans. Amer. Math. Soc., 362:10 (2010), pp. 5043–5062.
- [22] P. LI AND R. SCHOEN, *L^p and mean value properties of subharmonic functions on Riemannian manifolds*, Acta Math., 153:3-4 (1984), pp. 279–301.
- [23] P. LI AND J. WANG, *A generalization of Cheng's theorem*, Asian J. Math., 12:4 (2008), pp. 519–526.
- [24] ———, *Weighted Poincaré inequality and rigidity of complete manifolds*, Ann. Sci. École Norm. Sup. (4), 39:6 (2006), pp. 921–982. (English, with English and French summaries.)
- [25] J.-F. LI, *Eigenvalues and energy functionals with monotonicity formulas under Ricci flow*, Math. Ann., 338:4 (2007), pp. 927–946.
- [26] O. MUNTEANU, *Two results on the weighted Poincaré inequality on complete Kähler manifolds*, Math. Res. Lett., 14:6 (2007), pp. 995–1008.
- [27] L. E. PAYNE AND H. F. WEINBERGER, *An optimal Poincaré inequality for convex domains*, Arch. Rat. Mech. Anal., 5 (1960), pp. 286–292.
- [28] K. SEO, *Isoperimetric inequalities for submanifolds with bounded mean curvature*, Monatsh. Math., 166:3-4 (2012), pp. 525–542.
- [29] L. SIMON, *Lectures on geometric measure theory*, Proceedings of the Centre for Mathematical Analysis, Australian National University, Vol. 3, Australian National University, Centre for Mathematical Analysis, Canberra, 1983.
- [30] J. URBAS, *Monotonicity formulas and curvature equations*, J. Reine Angew. Math., 557 (2003), pp. 199–218.

